

Calculating the interaction force between the light beam and arbitrary-shape microparticles

R.V. Skidanov^{1,2}

¹ *Image Processing Systems Institute of RAS*

² *Samara State Aerospace University named after academician S.P. Korolev*

Abstract

The article describes a method for calculating the forces acting on a microparticle in a light field. The adequacy of the method is verified on the basis of comparison with experimental data.

Keywords: light beam, arbitrary-shape microparticles, microparticle.

Citation: Skidanov RV. Calculating the interaction force between the light beam and arbitrary-shape microparticles. *Computer Optics* 2005; 28: 18-21.

[Access full text \(in Russian\)](#)

References

- [1] Ashkin A, et al. Observation of a single-beam gradient force optical trap for dielectric particles. *Opt Lett* 1986; 11(5): 288-290.
- [2] Svoboda K, Block SM. Optical trapping of metallic Rayleigh particles. *Opt Lett* 1994; 19(13): 930-932.
- [3] Allen L, Eberly JH. *Optical resonance and two-level atoms*. New York: Dover Publications; 1987.
- [4] Kotlyar VV, Nalimov AG. Calculation of light pressure on a round dielectric cylinder using a fast iterative algorithm and based on an analytical solution. *Computer Optics* 2005; 27: 105-111.
- [5] Ashkin A, Dziedzic JM. Optical levitation by radiation pressure. *Appl Phys Lett* 1971; 19: 283.
- [6] Nieminen TA, Rubinsztein-Dunlop H, Heckenberg NR. Calculation and optical measurement of laser trapping forces on non-spherical particles. *J Quant Spectrosc Radiat Transf* 2001; 70: 627-637.